

Largemouth Bass (from NC WINS) ^[1]

Largemouth Bass

Micropterus salmoides

by Wayne Jones Updated by Kevin Dockendorf, Fisheries Biologist, 2009.
[N.C. Wildlife Resources Commission](#) ^[2].



Largemouth Bass

Classification

Class: Osteichthyes (bony fishes)
Order: Centrarchidae (sunfish)

Average Size

2 to 3 lbs.; the world record largemouth bass weighed 22 lbs, 4 oz. The North Carolina record is 15 lbs, 14 oz.

Food

Worms, salamanders, frogs, crayfish, snakes, ducklings, insects and fish.

Breeding

Spawning season occurs in spring when water temperatures reach 63 degrees.
Female deposits eggs in nest guarded by male bass. From 2,000 to 12,000 fry hatch. Only 5 to 10 will reach 10 in.

Young

When hatched, largemouth bass are known as fry. Fry are less than 1 inch long and eat zooplankton. As they grow from 1 to 3 inches, largemouth bass are known as fingerlings and eat aquatic insects and small fish.

Life Expectancy

Females live longer than males. In North Carolina, 10 years is long-lived. Most survive only 3 to 4 years.



The largemouth bass can be found in the
Range and Distributshaded regions. **ion**

In North Carolina, the largemouth bass is found in ponds, reservoirs, streams and rivers throughout the state, as well as in brackish coastal waters alongside saltwater species. Populations are considered good in every area of North Carolina, with the best populations occurring in the Piedmont and Coastal Plain. Bass tend to grow faster and larger in reservoirs, although large fish are also common in many farm ponds and in some coastal rivers.

General Information

Largemouth bass are the most sought-after freshwater gamefish in the United States. There are several reasons for the popularity of this fish. Anglers can usually find waters that contain largemouth bass within a short distance from their homes; they can catch bass that regularly reach trophy size (7 to 15 lbs.); and these fish are exciting to catch, with a battle characterized by head shaking and explosive leaps from the water.




Diagram of Largemouth Bass

Description

Largemouth bass belong to a group of fish collectively called black bass. Other black bass found in North Carolina include the smallmouth and spotted bass. All black bass belong to the sunfish family, but are distinguished from other sun fish by their elongated bodies. Largemouth bass are distinguished from other black bass by means of the lower jaw, which is longer than those of other bass. The largemouth bass is also a bigger species than the smallmouth or spotted bass. There are two subspecies of largemouth bass: the Florida largemouth bass and the northern largemouth bass.

History and Status

The largemouth bass is a native species of North Carolina. In North America it was originally found from southeastern Canada through the Great Lakes; south down the Mississippi Valley to Mexico and Florida; and up the Atlantic coast as far north as Maryland. As the fish grew in popularity with anglers, it was stocked in other areas and today is found throughout the continental United States and Hawaii, and southeastern Canada. It has also been stocked in Europe, Asia, Africa and South America.

Habitat and Habits

The largemouth bass prefers habitat with lots of structure. This structure may be in the form of weed beds, sunken logs, rocks, brush and standing timber. It uses structure as cover for ambush sites and to hide from larger predators.

Largemouth bass prefer temperatures between 77 and 86 degrees. Bass seldom feed at temperatures below 50 degrees and cannot survive for long at temperatures above 98 degrees.

When water temperatures approach 60 degrees in the spring, male bass swim to the spawning grounds. Male bass spend several days selecting their nest sites. The beds are usually in 1 to 4 ft. of water, but they may be deeper in clear water.

Spawning begins when the water reaches 63 to 68 degrees and remains in this temperature range for several days. A female largemouth bass lays between 2,000 and 7,000 eggs per pound of body weight.

The male guards the nest, fanning the eggs with his tail to keep off silt and debris. He will attack anything that swims near the nest. Despite the male's vigilance, many eggs are eaten by other fish species. Hatching time for bass eggs depends

on water temperatures. If the water is 72 degrees, eggs will hatch in only 2 days, but they take 5 days to hatch in water around 67 degrees. Approximately 2,000 to 12,000 fry hatch from each nest. Of these, however, only 5 to 10 are likely to survive to reach a size of 10 in. Newly hatched largemouth bass feed on tiny crustaceans and other zooplankton until they reach 2 in. in length. Young largemouth bass eat insects and small fish, including smaller bass. The adult largemouth bass will eat almost anything alive that will fit in their mouths. Fish, worms, frogs, insect larvae, crayfish, salamanders, snakes, small mammals, birds and ducklings are also eaten. Anglers use a variety of natural baits and artificial lures to catch largemouth bass.

Springtime movements center around spawning when bass move from deep to shallow water. During the summer months, bass are active in shallow waters during the morning and afternoons, but may move to deeper water at midday, when the water temperature increases in the shallows. When the water starts to cool in the fall, bass return to the shallows to look for food. As the water temperatures decrease during the winter months, bass become more sluggish, their metabolism slows down, and they require less food. They may become active during winter months if several days of



warm weather heat the water temperatures above 50 degrees. Largemouth bass fishing is a popular sport.

People Interactions

Largemouth bass have had a tremendous influence on many people. Tackle manufacturers, boat dealers and bait shops all provide specialized gear for largemouth bass fishing, especially for an increasing number of anglers involved in largemouth bass tournaments. Largemouth bass fishing is big business; many people make their livelihood from this popular sport. Although water pollution is a real threat to largemouth bass and other fish populations in some waters, the largemouth bass is an adaptable species and, overall, its future looks bright.



Biologists educate people about NCWRC Interaction Largemouth bass.

After fish kills associated with Hurricane Isabel occurred in northeast North Carolina in 2003, fisheries biologists began an intensive largemouth bass stocking project to evaluate stocking strategies that may be used after future hurricane-induced fish kills. In 2004, largemouth bass of different sizes (8, 2 and 5 inches, respectively) were cheek-tagged with a minute magnetic tag and stocked in the Roanoke and Chowan Rivers; stocking size was related to the time of year that certain sizes of largemouth bass may normally occur. Stockings were evaluated through an intensive electrofishing recapture period; the recapture period was completed in spring 2007. Although an initial boost of tagged individuals was observed in the population, the number of tagged largemouth bass contributing to the population over time dwindled to a mere fraction of the naturally produced largemouth bass. In this case, natural recruitment was likely a product of a few fish that must have survived to reproduce and repopulate a devastated fish community. Still, this study provided valuable insight on a management strategy to monitor largemouth bass recovery. Plans include implementing a sub-adult stocking strategy, when minimal to no natural recruitment occurs after hurricane-induced fish kills.

References:

- McClane, A.J. McClane's New Standard Fishing Encyclopedia (New York, N.Y.: Holt, Rinehart and Winston, 1965).
- Oster, Don. Largemouth Bass (W.A. Krueger Company, 1983).

Page, Lawrence M. and Burr, Brooks M. Peterson Field Guide to Freshwater Fishes (Boston, Mass.: Houghton Mifflin Company, 1991).

Pflieger, William I. The Fishes of Missouri (Missouri Department of Conservation, 1975).

Credits:

Illustrated by J.T. Newman. Photos by Dr. Amy Wethington, Chowan University and N.C. Wildlife Resources Commission.

Produced by the Division of Conservation Education, Cay Cross–Editor.



Profile courtesy of **NC WILDLIFE INFORMATION NETWORK SHARE (NC WINS)**

Subjects:

Wildlife ^[4]

Authors:

Dockendorf, Kevin ^[5]

Jones, Wayne ^[6]

Origin - location:

Coastal Plain ^[7]

Mountains ^[8]

Piedmont ^[9]

From:

NC Wildlife Information Network Share (NC WINS). ^[10]

11 June 2010 | Dockendorf, Kevin; Jones, Wayne

Source URL: <https://ncpedia.org/largemouth-bass-nc-wins?page=0>

Links

[1] <https://ncpedia.org/largemouth-bass-nc-wins> [2] <https://www.ncwildlife.org/> [3] <https://creativecommons.org/licenses/by-nd/4.0/> [4] <https://ncpedia.org/category/subjects/wildlife> [5] <https://ncpedia.org/category/authors/dockendorf-kevin> [6] <https://ncpedia.org/category/authors/jones-wayne> [7] <https://ncpedia.org/category/origin-location/coastal-pl> [8] <https://ncpedia.org/category/origin-location/mountains> [9] <https://ncpedia.org/category/origin-location/piedmont> [10] <https://ncpedia.org/category/entry-source/nc-wildlife-i>